

바젤 문제의 여러 가지 풀이

- 바젤 문제


$\sum_{n=1}^{\infty} \frac{1}{n^2}$ 을 닫힌 형식으로 구하여라!

* 닫힌 형식?

$$1 + \frac{1}{2^3} + \frac{1}{3^3} + \frac{1}{4^3} + \dots = 1.202 \dots \Rightarrow \text{닫힌 형식 } \times$$

$$1 + \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = 2 \Rightarrow \text{닫힌 형식 } \circ$$

풀이 ①: 2밀러의 풀이



$$\theta = \dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots$$

$$\Rightarrow \sin \theta = 0$$

$$\Rightarrow \sin x = 0 \text{ 일 때 } : \dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots$$

$$\sin x = kx \left(1 - \frac{x}{\pi}\right) \left(1 + \frac{x}{\pi}\right) \left(1 - \frac{x}{2\pi}\right) \left(1 + \frac{x}{2\pi}\right) \dots$$

($\because$ 바이어슈트라스의 곱정리)

$$\begin{aligned} \sin x &= kx \prod_{n=1}^{\infty} \left(1 - \frac{x}{n\pi}\right) \left(1 + \frac{x}{n\pi}\right) \\ &= kx \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right) \end{aligned}$$

$$\frac{\sin x}{x} = k \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right) \quad (x \neq 0)$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \lim_{x \rightarrow 0} k \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right)$$

$$1 = k \prod_{n=1}^{\infty} 1 = k \Rightarrow k=1$$

$$\frac{\sin x}{x} = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right)$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \dots$$

$$\frac{\sin x}{x} = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \frac{x^8}{9!} - \dots$$

$$\text{이차항계4: } -\frac{1}{3!} = -\frac{1}{6}$$

$$\frac{\sin x}{x} = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2\pi^2}\right)$$

$$= \left(1 - \frac{x^2}{\pi^2}\right) \left(1 - \frac{x^2}{2^2\pi^2}\right) \left(1 - \frac{x^2}{3^2\pi^2}\right) \dots$$

$\Rightarrow$ 모든 항이 상수와 이차항으로 이루어져 있음

$\Rightarrow$ 이차항: 모든 상수와 모든 이차항을 곱한 것들의 합

$$-\frac{x^2}{\pi^2} - \frac{x^2}{2^2\pi^2} - \frac{x^2}{3^2\pi^2} - \dots$$

$$\Rightarrow \text{이차항계4: } -\frac{1}{\pi^2} - \frac{1}{2^2\pi^2} - \frac{1}{3^2\pi^2} - \dots = -\frac{1}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

$$-\frac{1}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} = -\frac{1}{6}$$

$$\therefore \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

풀이 ②) 이 중적분 풀이

$$I_m = \int_1^0 x^m \ln x \, dx \quad (m=0, 1, 2, \dots)$$

$$I_0 = \int_1^0 \ln x \, dx = x(\ln x - 1) \Big|_1^0$$

$$= \lim_{b \rightarrow 0^+} x(\ln x - 1) \Big|_1^b = \lim_{b \rightarrow 0^+} \{b(\ln b - 1) + 1\}$$

$$= \lim_{b \rightarrow 0^+} (b \ln b - b + 1)$$

$$\left(\lim_{b \rightarrow 0^+} b \ln b = \lim_{b \rightarrow 0^+} \frac{\ln b}{\frac{1}{b}} = \lim_{b \rightarrow 0^+} \frac{\frac{1}{b}}{-\frac{1}{b^2}} = \lim_{b \rightarrow 0^+} -b = 0 \right)$$
$$= 0 - 0 + 1 = 1$$

$$\therefore I_0 = 1$$

$$I_m = \int_1^0 x^m \ln x \, dx \quad (m=1, 2, 3, \dots)$$

$$= \frac{x^{m+1}}{m+1} \ln x \Big|_1^0 - \int_1^0 \frac{x^m}{m+1} \, dx$$

$$= \lim_{b \rightarrow 0^+} \frac{x^{m+1} \ln x}{m+1} \Big|_1^b - \frac{x^{m+1}}{(m+1)^2} \Big|_1^0$$

$$= \lim_{b \rightarrow 0^+} \frac{b^{m+1} \ln b}{m+1} - \left(0 - \frac{1}{(m+1)^2} \right)$$

$$= \frac{1}{m+1} \lim_{b \rightarrow 0^+} \left\{ \underbrace{b^m}_{\downarrow 0} (\underbrace{b \ln b}_{\downarrow 0}) \right\} + \frac{1}{(m+1)^2} = \frac{1}{(m+1)^2}$$

$$\therefore I_m = \frac{1}{(m+1)^2} \quad (m=1, 2, 3, \dots)$$

$$I_0 = 1 \text{ 이므로 } I_m = \frac{1}{(m+1)^2} \quad (m=0, 1, 2, \dots)$$

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \sum_{n=1}^{\infty} \frac{1}{(2n)^2} + \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} + \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$$

$$\frac{3}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$$

$$\therefore \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{4}{3} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{4}{3} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^2} = \frac{4}{3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2}$$

$$= \frac{4}{3} \sum_{n=0}^{\infty} I_{2n}$$

$$\therefore I_m = \frac{1}{(m+1)^2}$$

$$= \frac{4}{3} \sum_{n=0}^{\infty} \int_1^0 x^{2n} \ln x \, dx$$

$$= \frac{4}{3} \int_1^0 \ln x \sum_{n=0}^{\infty} x^{2n} \, dx$$

$$= \frac{4}{3} \int_1^0 (\ln x) \frac{1}{1-x^2} \, dx \quad \because \sum_{n=0}^{\infty} x^{2n} = \frac{1}{1-x^2} \quad (0 < x < 1)$$

$$= \frac{4}{3} \int_1^0 \frac{\ln x}{1-x^2} dx$$

$$= \frac{4}{3} \int_1^0 \frac{1}{1-x^2} \left(\frac{1}{2} \ln(x^2) \right) dx$$

$$= \frac{4}{3} \cdot \frac{1}{2} \int_1^0 \frac{1}{1-x^2} (\ln x^2 - \ln 1) dx \quad \ln 1 = 0$$

$$\bullet \ln\left(\frac{1+x^2y^2}{1+y^2}\right) \quad y=0 \Rightarrow \ln 1$$

$$\lim_{n \rightarrow \infty} \ln\left(\frac{1+x^2y^2}{1+y^2}\right) = \lim_{n \rightarrow \infty} \ln\left(\frac{\frac{1}{y^2} + x^2}{\frac{1}{y^2} + 1}\right) = \ln x^2$$

$$\ln x^2 - \ln 1 = \ln\left(\frac{1+x^2y^2}{1+y^2}\right) \Big|_0^\infty$$

$$= \ln(1+x^2y^2) - \ln(1+y^2) \Big|_0^\infty$$

$$= \int_0^\infty \frac{2x^2y}{1+x^2y^2} - \frac{2y}{1+y^2} dy$$

$$= \int_0^\infty \frac{2x^2y(1+y^2) - 2y(1+x^2y^2)}{(1+x^2y^2)(1+y^2)} dy$$

$$2x^2y(1+y^2) - y(1+x^2y^2) = 2x^2y + 2x^2y^3 - 2y - 2x^2y^3$$

$$= 2x^2y - 2y$$

$$\int_0^{\infty} \frac{2y(x^2-1)}{(1+x^2y^2)(1+y^2)} dy$$

$$\frac{2}{3} \int_1^0 \frac{1}{1-x^2} (\ln x^2 - \ln 1) dx$$

$$= \frac{2}{3} \int_1^0 \frac{1}{1-x^2} \int_0^{\infty} \frac{2y(x^2-1)}{(1+x^2y^2)(1+y^2)} dy dx$$

$$= \frac{2}{3} \int_0^1 \frac{2(x^2-1)}{x^2-1} \int_0^{\infty} \frac{y}{(1+x^2y^2)(1+y^2)} dy dx$$

$$= \frac{4}{3} \int_0^1 \int_0^{\infty} \frac{y}{(1+x^2y^2)(1+y^2)} dy dx$$

$$= \frac{4}{3} \int_0^{\infty} \int_0^1 \frac{y}{(1+x^2y^2)(1+y^2)} dx dy$$

$$= \frac{4}{3} \int_0^{\infty} \frac{y}{1+y^2} \int_0^1 \frac{1}{1+x^2y^2} dx dy$$

$$\frac{d}{dx} \arctan x = \frac{1}{1+x^2}$$

$$\int_0^1 \frac{1}{1+y^2x^2} dx = \frac{1}{y} \arctan yx \Big|_0^1$$

$$= \frac{1}{y} \arctan y$$

$$\rightarrow = \frac{4}{3} \int_0^{\infty} \frac{y}{1+y^2} \cdot \frac{1}{y} \arctan y dy$$

$$= \frac{4}{3} \int_0^{\infty} \frac{\arctan y}{1+y^2} dy$$

$$= \frac{2}{3} \int_0^{\infty} 2\arctan y \cdot \frac{1}{1+y^2} dy$$

$$= \frac{2}{3} \left(\arctan y \right)^2 \Big|_0^{\infty}$$

$$\arctan y \xrightarrow{y \rightarrow \infty} \frac{\pi}{2}$$

$$= \frac{2}{3} \left\{ \left(\frac{\pi}{2} \right)^2 - 0 \right\} = \frac{2}{3} \cdot \frac{\pi^2}{4} = \frac{\pi^2}{6}$$

$$\therefore \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

$\frac{\pi}{2}$ | ③ : 코시의 증명

$$e^{i\theta} = \cos\theta + i\sin\theta$$

$$\theta = nx$$

$$e^{inx} = \cos nx + i\sin nx$$

$$e^{inx} = (e^{ix})^n = (\cos x + i\sin x)^n = \cos nx + i\sin nx$$

$$\frac{\cos(nx) + i\sin(nx)}{(\sin x)^n} = \frac{(\cos x + i\sin x)^n}{(\sin x)^n} = (\cot x + i)^n$$

$$(\cot x + i)^n = \sum_{r=0}^n \binom{n}{r} (\cot x)^{n-r} i^r$$

$$i^1 = i, i^2 = -1, i^3 = -i, i^4 = 1, i^5 = i, i^6 = -1, \dots$$

$$\sum_{r=0}^n \binom{n}{r} (\cot x)^{n-r} i^r = \underbrace{\sum_{r=0}^n \binom{n}{2r} (\cot x)^{n-2r} i^{2r}}_{\text{A4}} + \underbrace{\sum_{r=0}^n \binom{n}{2r+1} (\cot x)^{n-2r-1} i^{2r+1}}_{\text{B4}}$$

$$\sum_{r=0}^n \binom{n}{2r+1} (\cot x)^{n-2r-1} i^{2r+1}$$

$$= i \sum_{r=0}^n \binom{n}{2r+1} (\cot x)^{n-2r-1} i^{2r}$$

$$= i \sum_{r=0}^n \binom{n}{2r+1} (-1)^r (\cot x)^{n-2r-1} = i \frac{\sin(nx)}{(\sin x)^n}$$

$$\frac{\sin(nx)}{(\sin x)^n} = \sum_{r=0}^n \binom{n}{2r+1} (-1)^r (\cot x)^{n-2r-1}$$

$$n = 2m+1, x = u_k \quad \left(u_k = \frac{k\pi}{2m+1} \right) \quad (k=1, 2, 3, \dots, m)$$

$$nx = (2m+1) \cdot \frac{k\pi}{2m+1} = k\pi$$

$$\sin k\pi = 0 \quad \therefore \frac{\sin(nx)}{(\sin x)^n} = 0$$

$$0 = \sum_{r=0}^n \binom{2m+1}{2r+1} (-1)^r (\cot u_k)^{2m+1-2r-1}$$

$$= \binom{2m+1}{1} (\cot u_k)^{2m} - \binom{2m+1}{3} (\cot u_k)^{2m-2} + \binom{2m+1}{5} (\cot u_k)^{2m-4} - \dots$$

$$t_k = \cot^2 u_k$$

$$0 = \binom{2m+1}{1} t_k^m - \binom{2m+1}{3} t_k^{m-1} + \binom{2m+1}{5} t_k^{m-2} - \dots$$

↳ 해가 $t_1, t_2, t_3, \dots, t_m$ 일 가정

$$t_1 + t_2 + t_3 + \dots + t_m = \frac{\binom{2m+1}{3}}{\binom{2m+1}{1}} = \frac{(2m+1)2m(2m-1)}{3 \cdot 2 \cdot 1} = \frac{m(2m-1)}{3}$$


$$t_k = \cot^2 u_k$$

$$\sum_{k=1}^m t_k = \sum_{k=1}^m \cot^2 u_k = \frac{m(2m-1)}{3}$$

$$\cot^2 u_k + 1 = \csc^2 u_k \quad \cot^2 u_k = \csc^2 u_k - 1$$

$$\sum_{k=1}^m (\csc^2 u_k - 1) = \frac{m(2m-1)}{3} = \sum_{k=1}^m \csc^2 u_k - m$$

$$\sum_{k=1}^m \csc^2 u_k = \frac{m(2m-1)}{3} + m \quad \therefore \sum_{k=1}^m \csc^2 u_k = \frac{m(2m+2)}{3}$$



$$\sin x < x < \tan x$$

$$\cot x < \frac{1}{x} < \csc x$$

$$\cot^2 x < \frac{1}{x^2} < \csc^2 x \quad (\cot x > 0)$$

$$\sum_{k=1}^m \cot^2 u_k < \sum_{k=1}^m \frac{1}{u_k^2} < \sum_{k=1}^m \csc u_k^2$$

$$\frac{m(2m-1)}{3} < \sum_{k=1}^m \frac{1}{u_k^2} < \frac{m(2m+2)}{3}$$

$$u_k = \frac{k\pi}{2m+1}$$

$$\frac{m(2m-1)}{3} < \sum_{k=1}^m \frac{(2m+1)^2}{k^2 \pi^2} < \frac{m(2m+2)}{3}$$

$$\frac{\pi^2 m(2m-1)}{3(2m+1)^2} < \sum_{k=1}^m \frac{1}{k^2} < \frac{\pi^2 m(2m+2)}{3(2m+1)^2}$$

$$\lim_{m \rightarrow \infty} \frac{\pi^2 m(2m-1)}{3(2m+1)^2} = \lim_{m \rightarrow \infty} \frac{\pi^2}{3} \cdot \frac{2m^2 - m}{4m^2 + 4m + 1} = \frac{\pi^2}{3} \cdot \frac{1}{2} = \frac{\pi^2}{6}$$

$$\lim_{m \rightarrow \infty} \frac{\pi^2 m(2m+2)}{3(2m+1)^2} = \lim_{m \rightarrow \infty} \frac{\pi^2}{3} \cdot \frac{2m^2 + 2m}{4m^2 + 4m + 1} = \frac{\pi^2}{3} \cdot \frac{1}{2} = \frac{\pi^2}{6}$$

$$\frac{\pi^2}{6} \leq \lim_{m \rightarrow \infty} \sum_{k=1}^m \frac{1}{k^2} \leq \frac{\pi^2}{6}$$

$$\therefore \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$$

$\frac{\pi^2}{6}$ 이 ④ : 큰 그 함수의 이분

※ 소수와 원주율

- 에라토스테네스의 체

소수만 남기는 방법

	2	3	5	7	
11		13		17	19
		23			29
31				37	
41		43		47	
		53			59
61				67	
71		73			79
81		83			89
				97	

- 제타함수

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 1 + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \dots$$

우현에 1만 남겨보자!

$$\frac{1}{2^s} \zeta(s) = \frac{1}{2^s} + \frac{1}{4^s} + \frac{1}{6^s} + \frac{1}{8^s} + \dots$$

$$\zeta(s) - \frac{1}{2^s} \zeta(s) = 1 + \frac{1}{3^s} + \frac{1}{5^s} + \frac{1}{7^s} + \frac{1}{9^s} + \frac{1}{11^s} + \frac{1}{13^s} + \frac{1}{15^s} + \frac{1}{17^s} + \dots$$

$$= \left(1 - \frac{1}{2^s}\right) \zeta(s)$$

$$\frac{1}{3^s} \left(1 - \frac{1}{2^s}\right) \zeta(s) = \frac{1}{3^s} + \frac{1}{9^s} + \frac{1}{15^s} + \frac{1}{21^s} + \frac{1}{27^s} + \dots$$

$$\left(1 - \frac{1}{2^s}\right) \zeta(s) - \frac{1}{3^s} \left(1 - \frac{1}{2^s}\right) \zeta(s) = 1 + \frac{1}{5^s} + \frac{1}{7^s} + \frac{1}{11^s} + \frac{1}{13^s} + \frac{1}{17^s} + \dots$$

$$= \left(1 - \frac{1}{2^s}\right) \left(1 - \frac{1}{3^s}\right) \zeta(s)$$

⋮

$$\left(1 - \frac{1}{2^s}\right) \left(1 - \frac{1}{3^s}\right) \left(1 - \frac{1}{5^s}\right) \dots \zeta(s) = 1$$

$$\prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right) \zeta(s) = 1$$

$$\zeta(s) = \prod_{p \in \mathbb{P}} \left(1 - \frac{1}{p^s}\right)^{-1} = \prod_{p \in \mathbb{P}} (1 - p^{-s})^{-1}$$

$$\zeta(s) = \prod_{p \in \mathbb{P}} (1 - p^{-s})^{-1}$$

$$N = p_1^{a_1} p_2^{a_2} p_3^{a_3} p_4^{a_4} p_5^{a_5} \dots$$

$$p_1, p_2, p_3, \dots \in \mathbb{P} \quad (p_i < p_{i+1})$$

a_i 는 0이 아닌 정수 $\Rightarrow n$ 에 따라 변하는 4

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \sum_{a_1, a_2, a_3, \dots \geq 0} \frac{1}{p_1^{a_1 s} p_2^{a_2 s} p_3^{a_3 s} \dots}$$

$$= \sum_{a_1=0}^{\infty} \frac{1}{p_1^{a_1 s}} \sum_{a_2=0}^{\infty} \frac{1}{p_2^{a_2 s}} \sum_{a_3=0}^{\infty} \frac{1}{p_3^{a_3 s}} \dots$$

$$= \prod_{p \in \mathbb{P}} \sum_{a=0}^{\infty} \frac{1}{p^{a s}} = \prod_{p \in \mathbb{P}} \frac{1}{1 - \frac{1}{p^s}}$$

$$= \prod_{p \in \mathbb{P}} (1 - p^{-s})^{-1}$$

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \dots$$

$$= \prod_{p \in \mathbb{P}} (1 - p^{-2})^{-1} = \frac{\pi^2}{6}$$

$$= (1 - 2^{-2})^{-1} (1 - 3^{-2})^{-1} (1 - 5^{-2})^{-1} \dots$$

$$= \frac{1}{1 - \frac{1}{2^2}} \cdot \frac{1}{1 - \frac{1}{3^2}} \cdot \frac{1}{1 - \frac{1}{5^2}} \dots = \frac{\pi^2}{6}$$

$\frac{\pi^2}{6} \textcircled{4}$: 정수들의 역제곱의 합

$$\sin x = x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2 \pi^2}\right)$$

$$x = \pi y$$

$$\sin \pi y = \pi y \prod_{n=1}^{\infty} \left(1 - \frac{y^2}{n^2}\right)$$

$$\begin{aligned} \ln(\sin \pi y) &= \ln \pi y + \ln(1 - y^2) + \ln\left(1 - \frac{y^2}{2^2}\right) + \ln\left(1 - \frac{y^2}{3^2}\right) + \dots \\ &= \ln \pi y + \sum_{n=1}^{\infty} \ln\left(1 - \frac{y^2}{n^2}\right) \end{aligned}$$

$$\frac{\pi \cos \pi y}{\sin \pi y} = \frac{\pi}{\pi y} + \sum_{n=1}^{\infty} \frac{\frac{-2y}{n^2}}{1 - \frac{y^2}{n^2}}$$

$$= \frac{1}{y} + \sum_{n=1}^{\infty} \frac{-2y}{n^2 - y^2}$$

$$-\frac{\pi \cos \pi y}{2y \sin \pi y} = -\frac{1}{2y^2} + \sum_{n=1}^{\infty} \frac{1}{n^2 - y^2}$$

$$\frac{1}{2y^2} - \frac{\pi \cos \pi y}{2y \sin \pi y} = \sum_{n=1}^{\infty} \frac{1}{n^2 - y^2}$$

$$\lim_{y \rightarrow 0} \left(\frac{1}{2y^2} - \frac{\pi \cos \pi y}{2y \sin \pi y} \right) = \lim_{y \rightarrow 0} \sum_{n=1}^{\infty} \frac{1}{n^2 - y^2} = \sum_{n=1}^{\infty} \frac{1}{n^2}$$

$$\lim_{y \rightarrow 0} \left(\frac{1}{2y^2} - \frac{\pi \cos \pi y}{2y \sin \pi y} \right) = \lim_{y \rightarrow 0} \frac{\sin \pi y - \pi y \cos \pi y}{2y^2 \sin \pi y}$$

$$= \lim_{y \rightarrow 0} \frac{\pi \cos \pi y - \pi \cos \pi y + \pi^2 y \sin \pi y}{4y \sin \pi y + 2\pi y^2 \cos \pi y}$$

$$= \lim_{y \rightarrow 0} \frac{\pi^2 y \sin \pi y}{4y \sin \pi y + 2\pi y^2 \cos \pi y}$$

$$= \lim_{y \rightarrow 0} \frac{\pi^2 \sin \pi y}{4 \sin \pi y + 2\pi y \cos \pi y}$$

$$= \lim_{y \rightarrow 0} \frac{\pi^2}{4 + 2\pi y \frac{\cos \pi y}{\sin \pi y}}$$

$$\lim_{y \rightarrow 0} 2\pi y \frac{\cos \pi y}{\sin \pi y} = \lim_{y \rightarrow 0} 2\cos \pi y \cdot \frac{1}{\frac{\sin \pi y}{\pi y}} \rightarrow 1 = 2$$

$$\therefore \frac{\pi^2}{4+2} = \frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2}$$

//