

Higher homotopy group & Poincaré Conjecture

0. Prerequisites

The prerequisites for this seminar are the fundamental group, covering spaces, singular homology, cohomology, CW complex, Categorical Algebra.

1. Higher homotopy groups

(X, x_0) : pointed topological space

Definition 1.1

$\pi_k(X, x_0)$ is a set of homotopy class of maps $f: (I^k, \partial I^k) \rightarrow (X, x_0)$

equivalently, $f: (S^k, p) \rightarrow (X, x_0)$

With a group operation $*$: $\pi_k \times \pi_k \rightarrow \pi_k$: $[f] * [g] := [f * g]$

$$f * g := \begin{cases} f(2t_1, \dots) & t_1 \in [0, \frac{1}{2}] \\ g(2t_1 - 1, \dots) & t_1 \in [\frac{1}{2}, 1] \end{cases}$$

$\pi_k(X, x_0)$ is a group.

We already know that $\pi_1(X, x_0)$ can be nonabelian; figure-8.

However, $\pi_k(X, x_0)$ ($k \geq 2$) are always abelian.

Theorem 1.2 [Eckmann-Hilton]

$\times, \otimes$: unital binary operations on X .

Suppose $(a \times b) \otimes (c \times d) = (a \otimes c) \times (b \otimes d)$, $\forall a, b, c, d \in X$.

Then, $\times, \otimes$ are exactly same operation and commutative & associative.

Proof) Left to a reader except units.

$1_\times, 1_\otimes$: units

$$\begin{aligned} 1_\times &= 1_\times \times 1_\times = (1_\times \otimes 1_\otimes) \times (1_\otimes \otimes 1_\times) = (1_\times \times 1_\otimes) \otimes (1_\otimes \times 1_\times) \\ &= 1_\otimes \otimes 1_\otimes = 1_\otimes \end{aligned}$$

Definition 1.3

$$\otimes : \Pi_k \times \Pi_k \rightarrow \Pi_k : [f] \otimes [g] := [f \otimes g]$$

$$f \otimes g := \begin{cases} f(t_1, 2t_2, \dots) & t_2 \in [0, \frac{1}{2}] \\ g(t_1, 2t_2 - 1, \dots) & t_2 \in [\frac{1}{2}, 1] \end{cases}$$

Theorem 1.4

$(\pi_k, *)$ is an abelian group.

Proof) $\forall f, g, h, k : (S^k, p) \rightarrow (X, x_0)$, $(f \otimes g) * (h \otimes k) = (f * h) \otimes (g * k)$

$$\therefore ([f] \otimes [g]) * ([h] \otimes [k]) = ([f] * [h]) \otimes ([g] * [k])$$

By Eckmann-Hilton argument, $*$ is abelian, as desired.

Theorem 1.1 [Lifting Criterion]

If $p: \tilde{X} \rightarrow X$ is a covering map, $f: (Y, y_0) \rightarrow (X, x_0)$, $p(\tilde{x}_0) = x_0$, then f lifts to a $\tilde{f}: (Y, y_0) \rightarrow (\tilde{X}, \tilde{x}_0)$ with $p \circ \tilde{f} = f$ if and only if

$$f_* (\pi_1(Y, y_0)) \subseteq p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

$$\begin{array}{ccc} & (\tilde{X}, \tilde{x}_0) & \\ \tilde{f} \nearrow & \downarrow p & \\ (Y, y_0) & \xrightarrow{f} & (X, x_0) \end{array} \iff f_* (\pi_1(Y, y_0)) \subseteq p_* (\pi_1(\tilde{X}, \tilde{x}_0))$$

Theorem 1.6

If $k \geq 2$, $\pi_k(S^1, p) = 0$

Proof) Recall $p: \mathbb{R} \rightarrow S^1$: covering map. $\forall f: S^k \rightarrow S^1$ lifts to a map $\tilde{f}: S^k \rightarrow \mathbb{R}$, by the Lifting Criterion.

Since $\mathbb{R}$ is contractible, $\tilde{f}$ is nullhomotopic, i.e., f is nullhomotopic.

2. Hurewicz Isomorphism theorem.

Definition 2.1 [Hurewicz homomorphism]

$$\Phi: \pi_k(X, x_0) \rightarrow H_k(X)$$

$$[f] \mapsto f_*[S^k] \in H_k(X)$$

Since homology is homotopy invariance, Φ is well-defined.

Please check that Φ is a homomorphism.

Lemma 2.2

(X, x_0) : pointed space

i) $\forall k \geq 1$, if $f: (S^k, p) \rightarrow (X, x_0)$ is nullhomotopic without fixing the base points, then $[f] = 0 \in \pi_k(X, x_0)$

ii) $\forall k \geq 2$, $f: \partial \Delta_{k+1} \rightarrow X$: map sending $\forall (k-1)$ -dimensional faces to x_0 .

$$\text{Then, } [f] = \sum_{\sigma \in F_k \Delta_{k+1}}^* [f|_{\sigma}] \in \pi_k(X, x_0)$$

Sketch of Proof) i) Given a homotopy $F: S^k \times I \rightarrow X$

$$F(s, 0) = f$$


$$F(s, 1) = \text{const.}$$

We can use $\gamma: I \rightarrow X$: trajectory of p to modify F to homotopy
 $t \mapsto F(p, t)$

sending p to x_0 at all times.

(i) [Draw Picture]

$k=2$



Theorem 2.3 [Hurewicz isomorphism theorem]

$k \geq 2$, X : path-connected & $\pi_i(X, x_0) = 0, \forall i < k$.

Then Φ is an isomorphism, i.e., $\pi_k(X, x_0) \simeq H_k(X)$


Proof) Using Singular homology, define $\Psi: C_k(X) \rightarrow \pi_k(X, x_0)$

The idea is modify image of boundary via chain homotopy.

Define $K: C_i(X) \rightarrow C_{i+1}(X), \forall 0 \leq i < k$ as follows.

Since X : path-connected, $\forall p \in X$, choose a path $K(p)$ from x_0 to p ; $K(p): \Delta_1 \rightarrow X$ $K(p)(0) = x_0, K(p)(1) = p$

For each 1-simplex $\sigma: \Delta_1 \rightarrow X$, $\exists \tau: \partial\Delta_2 \rightarrow X$ sending the three faces to $\sigma, K(\sigma(0)), K(\sigma(1))$.



Since X is simply connected, τ can be extended to

$$K(\sigma): \Delta_2 \rightarrow X \text{ such that } \partial K(\sigma) + K(\partial\sigma) = \sigma$$

Continuing by induction on τ , if $1 \leq \tau < k$, then $\forall$ τ -simplex

$\sigma: \Delta_\tau \rightarrow X$, choose $(\tau+1)$ -simplex $K(\sigma): \Delta_{\tau+1} \rightarrow X$ which sends the faces to σ , and the faces of $K(\partial\sigma)$ and therefore satisfies

$$\partial K(\sigma) + K(\partial\sigma) = \sigma$$

Finally if $\sigma: \Delta_k \rightarrow X$ is a k -simplex, then $\exists \tau: \partial\Delta_{k+1} \rightarrow X$ sending the faces to σ , and the faces of $K(\partial\sigma)$.

Let q denote the face sent to x_0 .

Identifying $(\partial\Delta_{k+1}, q) \simeq (S^k, p)$ this gives $\underline{\Psi}(\sigma): (S^k, x_0) \rightarrow (X, x_0) \in \pi_k(X, x_0)$

Moreover, the sum of the faces is homologous to $\underline{\Psi}(\sigma)$ regarded as a simplex, i.e., $\exists$ a simplex $K(\sigma)$ with

$$\partial K(\sigma) + K(\partial\sigma) = \sigma - \Phi(\underline{\Psi}(\sigma))$$

While $\underline{\Psi}$ may depend on the above choices, we claim that

$\underline{\Psi}$ induces a map on homology which is inverse to Φ .

Theorem 2.4

$$\pi_k(S^n) = \begin{cases} 0 & \text{if } k < n \\ \mathbb{Z} & \text{if } k = n \end{cases}$$

Proof) By the Hurewicz isomorphism theorem, it's clear.

3. Dependence of π_k on the base point

We will show that if X is path connected, then the homotopy groups of X for different choices of base point are isomorphic.

If $\gamma: I \rightarrow X$: path from x_0 to x_1 , define a map



$$\Phi_\gamma: \pi_k(X, x_1) \rightarrow \pi_k(X, x_0) \text{ as follows}$$

Let us temporarily reparametrize $I^k = [-1, 1]^k$

If $f: [-1, 1]^k \rightarrow X$ & $t = (t_1, \dots, t_k) \in I^k$

let $m = \max\{|t_i|\}$ and define

$$\Phi_\gamma(f(t)) := \begin{cases} f(2t) & m \leq \frac{1}{2} \\ \gamma(2(1-m)) & m \geq \frac{1}{2} \end{cases}$$



Please check that Φ_γ is group homomorphism.

The following facts are easy to see

(i) If $\gamma \simeq_h \gamma'$, then $\Phi_\gamma = \Phi_{\gamma'} : \pi_k(X, x_0) \rightarrow \pi_k(X, x_1)$

(ii) If γ : path from x_0 to x_1 , γ' : path from x_1 to x_2 ,
then $\Phi_{\gamma_2 * \gamma_1} = \Phi_{\gamma_2} \circ \Phi_{\gamma_1} : \pi_k(X, x_2) \rightarrow \pi_k(X, x_0)$

(iii) If γ : constant path at x_0 , then $\Phi_\gamma = \bar{i}_{\pi_k(X, x_0)}$

Theorem 3.1

If X : path connected, then $\pi_k(X, x_0) \simeq \pi_k(X, x_1)$

4. Poincaré conjecture

Theorem 4.1. [Poincaré duality]

If M is n -dimensional orientable closed manifold, then

$$H_{n-k}(M) \simeq H^k(M)$$

Definition 4.2 [Hom, Ext Functor]

$A, B : R$ -module

$\text{Hom}_R(A, B) := \{ \text{Homomorphism from } A \text{ to } B \}$

For a fixed A , Hom_R is left exact functor.

For a injective resolution, $0 \rightarrow B \rightarrow I^0 \rightarrow I^1 \rightarrow \dots$

Form the cochain complex, $0 \rightarrow \text{Hom}_R(A, I^0) \rightarrow \text{Hom}_R(A, I^1) \rightarrow$

$\text{Ext}_R^i(A, B) := i\text{th cohomology of this complex}$

Ext_R^i is right derived functor from Hom_R

Theorem 4.3 [Universal coefficient theorem for cohomology]

R : PID, G : R -module, $\exists$ short exact sequence

$$0 \rightarrow \text{Ext}_R^1(H_{i-1}(X; R), G) \rightarrow H^i(X; G) \rightarrow \text{Hom}_R(H_i(X; R), G) \rightarrow 0$$

and the sequence splits.

Theorem 4.4 [Homology Whitehead theorem]

X, Y : simply connected CW complexes

$$f: X \rightarrow Y.$$

If $f_*: H_n(X) \rightarrow H_n(Y)$ is isomorphism $\forall n$,

then f : homotopy equivalence.

Theorem 4.5

Every simply connected, orientable, closed 3-manifold is homotopy equivalent to S^3 .

Proof) X : simply connected, orientable, closed 3-manifold

Then, $H_0(X) \cong \mathbb{Z}$, $H_1(X) = \pi_1(X) = 0$

Since X is orientable, $H_3(X) \cong \mathbb{Z}$.

By Poincaré duality, $H_2(X) \cong H^1(X)$

By Universal coefficient theorem,

$$H^1(X) \cong \text{Hom}(H_1(X); \mathbb{Z}) \oplus \text{Ext}(H_0(X); \mathbb{Z}) = 0$$

So $H_2(X) = 0$.

By Hurewicz theorem, $\Phi: \pi_3(X) \rightarrow H_3(X)$

is an isomorphism

Let $f: S^3 \rightarrow X$ represent a generator of $\pi_3(X)$

Then, $f_* : H_n(S^3) \rightarrow H_n(X)$ are isomorphisms

It follows that f is a homotopy equivalence

Theorem 4.6 [Poincaré Theorem]

Such X is, in fact, homeomorphic to S^3 .