

11.27. MIMIC Seminar.

< Differential Forms & de Rham Cohomology >.

1. Review on Calculus.

- (Fundamental theorem of Line integral)

$$f: U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}, \quad \gamma: [a, b] \rightarrow U \text{ smooth.}$$

$$\Rightarrow \int_{\gamma} \text{grad } f \cdot ds = f(\gamma(b)) - f(\gamma(a)) = \int_{\partial[a, b]} f.$$

- (Green's theorem)

$$D \subseteq \mathbb{R}^2, \quad F: \text{smooth v.f. on } D$$

$$\int_{\partial D} F \cdot ds = \iint_D \text{rot } F \, dA$$

- (Divergence Theorem)

$$R \subset \mathbb{R}^3, \quad F: \text{smooth v.f. on } R,$$

$$\iint_{\partial R} F \cdot dS = \iiint_R \text{div } F \, dV.$$


- (Stokes' theorem)

$$F: \text{smooth v.f. on a surface } S \subseteq \mathbb{R}^3.$$

$$\iint_S \text{curl } F \cdot dS = \int_{\partial S} F \cdot ds.$$

* Geometric Tangent Spaces.

$$\begin{aligned}\mathbb{R}_p^n &:= \text{the tangent space of } \mathbb{R}^n \text{ at } p. \\ &= \{p\} \times \mathbb{R}^n = \{(p, v) : v \in \mathbb{R}^n\}.\end{aligned}$$


$$\begin{array}{c} \nearrow v \\ \bullet \\ p \end{array} \rightsquigarrow \text{We abbreviate } (p, v) \text{ as } v_p.$$

$T_p \mathbb{R}^n$ forms a real vector space under the canonical operations :

$$v_p + w_p = (v + w)_p, \quad c \cdot v_p = (cv)_p.$$

For a smooth function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we define the directional derivative in the direction v at p by

$$D_v f(p) := \left. \frac{d}{dt} \right|_{t=0} f(p + tv)$$

$\Rightarrow$ We use the notation $D_v|_p f$ instead of $D_v f(p)$

$\Rightarrow D_v|_p f$ can be considered as

$$D_v|_p : C^\infty(\mathbb{R}^n) \rightarrow \mathbb{R} \quad (\text{called a derivation})$$

satisfying the following properties :

- $D_v|_p$ is linear.

- $D_v|_p(fg) = D_v|_p(f)g(p) + f(p)D_v|_p(g)$. (Leibniz Rule)

Let $T_p \mathbb{R}^n = \{ \text{all derivations of } C^\infty(\mathbb{R}^n) \text{ at } p \}$.

$\Rightarrow T_p \mathbb{R}^n$ forms a vector space under the operations:

$$(v+w)f = vf + wf, \quad (cv)f = c(vf)$$

for $f \in C^\infty(\mathbb{R}^n)$

One can check $\mathbb{R}_p^n \longrightarrow T_p \mathbb{R}^n$ is an isomorphism.

$$v_p \longmapsto D_v|_p$$

$$\{ \text{Tangent vectors on } \mathbb{R}^n \} = \{ \text{Derivations on } C^\infty(\mathbb{R}^n) \}.$$

In this reason,

$$D_{e_i}|_p f = \lim_{t \rightarrow 0} \frac{f(p+te_i) - f(p)}{t} = \frac{\partial f}{\partial x_i}(p) := \frac{\partial}{\partial x_i}|_p f$$

$\rightsquigarrow \{ \frac{\partial}{\partial x_i}|_p \}$ is a basis for $T_p \mathbb{R}^n$.

$\rightsquigarrow$ For fixed $f \in C^\infty(\mathbb{R}^n)$,

$$D_v|_p f = df_p(v) \rightsquigarrow df_p: T_p \mathbb{R}^n \rightarrow \mathbb{R} \text{ linear.}$$

$\Rightarrow df_p \in (T_p \mathbb{R}^n)^* = T_p^* \mathbb{R}^n$. : the cotangent space of $\mathbb{R}^n$ at p .

$\rightsquigarrow \{ dx_1|_p, \dots, dx_n|_p \}$ is a basis for $T_p^* \mathbb{R}^n$.

$\uparrow$ dual basis of $\{ \frac{\partial}{\partial x_1}|_p, \dots, \frac{\partial}{\partial x_n}|_p \}$.

2. Differential Forms on $\mathbb{R}^n$.

→ $\mathbb{R}$ -v.s + ring structure.

Define Ω^* to be the $\mathbb{R}$ -algebra generated by $dx_1, \dots, dx_n$ where $x_1, \dots, x_n$ are the standard linear coordinates on $\mathbb{R}^n$ with the relation

$$\bullet \quad dx_i dx_i = 0 \quad \forall i = 1, 2, \dots, n.$$

$$\bullet \quad dx_i dx_j = -dx_j dx_i \quad \forall i \neq j$$

⇒ Ω^* has a basis

$$1, dx_i, dx_i dx_j, dx_i dx_j dx_k, \dots, dx_1 \dots dx_n$$

$i < j \qquad i < j < k$

The (smooth) differential forms on $\mathbb{R}^n$ are elements of

$$\Omega^*(\mathbb{R}^n) = C^\infty(\mathbb{R}^n) \otimes_{\mathbb{R}} \Omega^*$$

(i.e. if $\omega \in \Omega^*(\mathbb{R}^n)$, ω can be uniquely written as

$$\omega = \sum_{i_1 < \dots < i_p} f_{i_1 i_2 \dots i_p} dx_{i_1} \dots dx_{i_p} = \sum f_I dx_I. \quad \leadsto p\text{-form}$$

$$\Rightarrow \Omega^*(\mathbb{R}^n) = \bigoplus_{p=0}^n \Omega^p(\mathbb{R}^n) \quad \text{where } \Omega^p(\mathbb{R}^n) = \{p\text{-forms on } \mathbb{R}^n\}.$$

is graded $\mathbb{R}$ -algebra.

3. Exterior Derivatives & de Rham Complex.

We have seen that for $f \in C^\infty(\mathbb{R}^n)$,

$$df_p : T_p \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\begin{array}{ccc} v = \sum_{i=1}^n v_i \frac{\partial}{\partial x_i} \Big|_p & \longmapsto & \sum_{i=1}^n v_i \frac{\partial f}{\partial x_i}(p) \\ \parallel & & \parallel \\ (v_1, \dots, v_n) & \longmapsto & v \cdot \text{grad} f(p) \end{array}$$

$$\Rightarrow \text{Since } dx_i|_p(v) = dx_i|_p\left(\sum_{j=1}^n v_j \frac{\partial}{\partial x_j} \Big|_p\right) = \sum_{j=1}^n v_j \delta_{ij} = v_i$$

We can write

$$df_p(v) = \sum_{i=1}^n v_i \frac{\partial f}{\partial x_i}(p) = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i|_p(v)$$

$$\Rightarrow df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i.$$

$\leadsto$ Generalization : There is a differential operator

$$d : \Omega^p(\mathbb{R}^n) \rightarrow \Omega^{p+1}(\mathbb{R}^n)$$

defined as follows :

① if $f \in \Omega^0(\mathbb{R}^n) \cong C^\infty(\mathbb{R}^n)$, $df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i$.

② if $\omega = \sum f_I dx_I$, $d\omega = \sum df_I dx_I$.

d is called the exterior differentiation.

The wedge product of $\omega = \sum f_I dx_I$, $\tau = \sum g_J dx_J$

$$\leadsto \omega \wedge \tau = \sum f_I g_J dx_I dx_J.$$

Properties

$$\text{i)} \quad \omega \wedge \tau = (-1)^{\deg \omega \deg \tau} \tau \wedge \omega. \quad (\text{graded-commutative})$$

$$\text{ii)} \quad d(\omega \wedge \tau) = d\omega \wedge \tau + (-1)^{\deg \omega} \omega \wedge d\tau \quad (d: \text{antiderivation})$$

$$\text{iii)} \quad d^2 = 0.$$

Proof of (iii) for $f \in \Omega^0(\mathbb{R}^n)$,

$$d^2 f = d\left(\sum_j \frac{\partial f}{\partial x_j} dx_j\right) = \sum_{i,j} \frac{\partial^2 f}{\partial x_i \partial x_j} dx_i dx_j = 0.$$

$$\text{For } \omega \in \Omega^p(\mathbb{R}^n), \quad \omega = \sum f_I dx_I$$

$$\begin{aligned} d^2 \omega &= d\left(\sum df_I dx_I\right) = \sum (d^2 f_I \wedge dx_I - df_I \wedge d^2 x_I) \\ &= 0. \end{aligned}$$

$$\dots \rightarrow \Omega^{p-1}(\mathbb{R}^n) \xrightarrow{d} \Omega^p(\mathbb{R}^n) \xrightarrow{d} \Omega^{p+1}(\mathbb{R}^n) \rightarrow \dots$$

: de Rham complex.

kernel of d = closed forms.

Image of d = exact forms.

Note: Since $d^2 = 0$, the exact forms are closed

Remark

The de Rham complex can be viewed as a set of differential equation whose solutions are closed forms.

For example, the differential equation $\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} = 0$ corresponds to the finding a closed 1-form $f dx + g dy$.

Remark On $\mathbb{R}^3$, There is the correspondence

$$\begin{array}{ccccccc} C^\infty(\mathbb{R}^3) & \xrightarrow{\text{grad}} & \mathcal{X}(M) & \xrightarrow{\text{curl}} & \mathcal{X}(M) & \xrightarrow{\text{div}} & C^\infty(\mathbb{R}^3) \\ \textcircled{1} \parallel & & \textcircled{2} \parallel & & \textcircled{3} \parallel & & \textcircled{4} \parallel \\ \Omega^0(\mathbb{R}^3) & \xrightarrow{d} & \Omega^1(\mathbb{R}^3) & \xrightarrow{d} & \Omega^2(\mathbb{R}^3) & \xrightarrow{d} & \Omega^3(\mathbb{R}^3) \end{array}$$

$$\textcircled{1} : f \longleftrightarrow f.$$

$$\textcircled{2} : X = (f_1, f_2, f_3) \longleftrightarrow f_1 dx + f_2 dy + f_3 dz.$$

$$\textcircled{3} : X = (f_1, f_2, f_3) \longleftrightarrow f_1 dy dz - f_2 dx dz + f_3 dx dy.$$

$$\textcircled{4} : f \longleftrightarrow f dx dy dz.$$

$$\begin{aligned} * \quad \text{curl}(\text{grad } f) &= 0 \\ \text{div}(\text{curl } F) &= 0 \end{aligned} \longleftrightarrow d^2 = 0.$$

$$\int_M d\omega = \int_{\partial M} \omega \quad (\text{Stokes' theorem}).$$

Definition The p -th de Rham cohomology $\checkmark$ of $\mathbb{R}^n$ is the quotient vector space defined by

$$H_{\text{dR}}^p(\mathbb{R}^n) = \frac{\{\text{closed } p\text{-forms}\}}{\{\text{exact } p\text{-forms}\}}.$$

* We can define $H_{dR}^p(U)$ for an open set $U \subseteq \mathbb{R}^n$ in the same way.

(i.e. $\Omega^*(U) = C^\infty(U) \otimes_{\mathbb{R}} \Omega^*$.)

Example.

$$(i) \quad H_{dR}^p(*) = \begin{cases} \mathbb{R} & p=0 \\ 0 & 0 < p \end{cases}$$

$$0 \xrightarrow{0} \Omega^0(*) \xrightarrow{0} 0$$

$$\quad \quad \quad \parallel$$

$$\quad \quad \quad \mathbb{R}$$

$$(ii) \quad H_{dR}^p(\mathbb{R}) = \begin{cases} \mathbb{R} & p=0 \\ 0 & 0 < p \end{cases}$$

$$0 \rightarrow \Omega^0(\mathbb{R}) \xrightarrow{d} \Omega^1(\mathbb{R}) \rightarrow 0$$

$$\Rightarrow \ker d \cap \Omega^0(\mathbb{R}) = \{\text{constant functions on } \mathbb{R}\} = \mathbb{R}.$$

$$\Rightarrow H_{dR}^0(\mathbb{R}) = \mathbb{R}.$$

For $H_{dR}^1(\mathbb{R})$, $\ker d = \Omega^1(\mathbb{R})$.

Let $\omega = f(x)dx$. consider $g(x) = \int_0^x f(t)dt$

$$\Rightarrow dg = f(x)dx = \omega. \Rightarrow H_{dR}^1(\mathbb{R}) = 0.$$

(iii) In general, $H_{dR}^p(\mathbb{R}^n) = \begin{cases} \mathbb{R} & p=0 \\ 0 & 0 < p \end{cases}$ (Poincaré lemma)

$$(iv) \quad H_{dR}^p(\mathbb{R} - \{0\}) = \begin{cases} \mathbb{R} \oplus \mathbb{R} & p=0 \\ 0 & 0 < p \end{cases} \rightsquigarrow \int \frac{1}{x} dx = \begin{cases} \log|x| + C_1 & (x > 0) \\ \log|x| + C_2 & (x < 0) \end{cases}$$

4. Functoriality.

Let $F: M \rightarrow N$ be a smooth map. for $\eta \in \Omega^p(N)$,
 $(F^*\eta)_p = dF_p^*(\eta_{F(p)})$

The pullback $F^*: \Omega^*(N) \rightarrow \Omega^*(M)$ carries

closed, exact forms on N into closed, exact forms on M .

And hence, it induces

$$F^*: H_{dR}^*(N) \rightarrow H_{dR}^*(M).$$

$\leadsto H_{dR}^*: \text{Man}^\infty \rightarrow \text{Vect}_\mathbb{R}$ defines a contravariant functor.


Example $H_{dR}^*(S^1)$

$$0 \xrightarrow{0} \Omega^0(S^1) \xrightarrow{d} \Omega^1(S^1) \xrightarrow{0} 0$$

$$H_{dR}^0(S^1) = \{ \text{closed 0-forms on } S^1 \} / \{0\}$$

$$= \{ f: S^1 \rightarrow \mathbb{R} \mid df = 0 \}$$

$S^1 \subset \mathbb{R}^2$, consider a coordinate $(\cos \theta, \sin \theta) \xrightarrow{\varphi} \theta$



$$df = \frac{df}{d\theta} d\theta = 0 \Leftrightarrow \frac{df}{d\theta} \equiv 0.$$

$\Rightarrow f$ is constant since S^1 is connected.

$$\Rightarrow H_{dR}^0(S^1) = \mathbb{R}.$$

$$H_{\text{dR}}^1(S^1) = \frac{\{\text{closed 1-forms on } S^1\}}{\{\text{exact 1-forms on } S^1\}}$$

Note that $\{\text{closed 1-forms on } S^1\} = \Omega^1(S^1)$.

Let $\xi \in \Omega^1(S^1)$, write $\xi = f(t)dt$.

If ξ is exact, i.e. $\xi = df$ for some $f \in \Omega^0(S^1)$

$$\int_{S^1} \xi = \int_{S^1} df = f(2\pi) - f(0) = 0. \quad \nearrow \int_{S^1} : \Omega^1(S^1) \rightarrow \mathbb{R} \text{ linear map.}$$

$$\Rightarrow \{\text{exact 1-forms on } S^1\} \subseteq \ker \int_{S^1}$$

If $\eta \in \Omega^1(S^1)$ satisfies $\int_{S^1} \eta = 0$

Write $\eta = f(t)dt$, define $g(t) = \int_0^t f(s)ds$

$$\Rightarrow dg = f(t)dt = \eta.$$

$$\therefore \ker \int_{S^1} = \{\text{exact 1-forms on } S^1\}$$

Hence, Isomorphism theorem says

$$H_{\text{dR}}^1(S^1) = \frac{\Omega^1(S^1)}{\ker \int_{S^1}} \cong \text{Im } \int_{S^1} = \mathbb{R}.$$

$$(\because \int_{S^1} \frac{dt}{2\pi} = 1)$$

